

Pluri-polarity in almost complex structures.

Jean-Pierre Rosay

arXiv:0902.4452v1 [math.CV] 25 Feb 2009

Abstract: J -holomorphic curves are $-\infty$ sets of J -plurisubharmonic functions, with a singularity of LogLog type, but it is shown that in general they are not $-\infty$ sets of J -plurisubharmonic functions with Logarithmic singularity (i.e. non-zero Lelong number). Some few additional remarks on pluripolarity in almost complex structures are made.

A.M.S. Classification: 32Q60, 32Q65, 32U05

INTRODUCTION. J -plurisubharmonic functions with poles (at which the function is $-\infty$) have already played a role in almost complex analysis. They have been used in the study of the Kobayashi metric ([G-S] [I-R]) for getting an efficient control of J -holomorphic discs. In a work in progress with S. Ivashkovich, applications to uniqueness problems are given. The first pluripolarity result is due to Chirka who showed that if J is a $\mathcal{C}^1$ almost complex structure defined near 0 in $\mathbf{C}^n$ and if $J(0) = J_{st}$ (the standard complex structure), then for $A > 0$ large enough $\log|z| + A|z|$ is J -plurisubharmonic near 0. A complete proof has been written in [I-R] (Lemma 1.4 page 2400). The function $-\log|\log|z||$ is also J -plurisubharmonic near 0. Although a ‘log-log singularity’ (zero Lelong number!) is much less interesting and has less applications than a ‘log singularity’, functions with log-log singularity were introduced in [Ro1] to show pluripolarity of J -holomorphic curves. Later, Elkhadhra [E] generalized the result to show pluri-polarity of J -holomorphic submanifolds, again with a ‘loglog’ singularity of the function.

In [Ro1] there has been an error in stating the smoothness hypotheses, all smoothness requirements in the statements have to be increased by 1 ($\mathcal{C}^{k+1}$ instead of $\mathcal{C}^k$). Indeed on line 11- page 663 it is claimed that because $J[Y', JY'](Z', 0) = 0$, one has $|J[Y', JY'](Z', Z'')| \leq C|Z''||Y'|^2$. This is correct if J is of class $\mathcal{C}^2$ (or at least $\mathcal{C}^{1,1}$), but $\mathcal{C}^1$ smoothness, hence continuity of $[Y', JY']$, is not enough.

The really original part of the paper is an example in dimension 4 showing that *in general, J -holomorphic curves are not $-\infty$ set of any J -plurisubharmonic function with a logarithmic singularity*. The precise statement is given in Proposition 2.

The plan of the paper is as follows:

Part 0 is a naive discussion, in one complex variable, of the functions $\log|z|$ and $\log|\log|z||$, and of perturbations of the Laplacian. Since the theory of almost complex structures is in part the theory of small perturbations of the standard complex structure (and thus of the Laplacian), it seems to me that this preliminary discussion leads to a better understanding of the problems that are encountered.

In Part I, we develop the calculus tools to be used in part II and III, for studying J -plurisubharmonicity.

Part II is a proof of the pluripolarity of J -holomorphic curves established in [Ro1] (with the smoothness hypothesis this time correctly stated!). Although the result is not new, the proof is written very differently, in a way such that I think that it is much easier to see how things happen. For me, it has been a needed preliminary before getting the counterexample that is given in Part III.

Part III itself consists of two parts. In the first one one shows the non existence of functions with log singularity under natural reasonable hypotheses on the behavior of the function. Then in the second part the non existence result is proved in full generality. For that, we shall need 3 Lemmas in measure theory whose proof is given in an Appendix.

Part IV treats the case when there is better matching with the standard complex structure, when the Nijenhuis tensor vanishes along a J -holomorphic disc. Things become much easier and functions with logarithmic poles exist.

Part V is devoted to the very simple remark that in dimension 4 a 2 dimensional surface cannot be pluripolar if it is not a J -holomorphic curve. In higher dimensions $2n$, the corresponding simple remark is for generic n -dimensional manifolds. Our elementary argument avoids any construction of 'Pinchuck discs' with part of their boundary attached to the submanifold.

We shall take the following definition of J -plurisubharmonicity:

Definition. *Let J be an almost complex structure defined on some smooth manifold, of class at least $\mathcal{C}^{1,\alpha}$ for some $0 < \alpha < 1$. An upper semi-continuous (hence continuous) function λ , defined on that set, with values in $[-\infty, +\infty)$ will be said to be J -plurisubharmonic if λ is of class $\mathcal{C}^2$ off its $-\infty$ set and if its restriction to any J -holomorphic curve is subharmonic.*

The hypothesis that J is of class $\mathcal{C}^{1,\alpha}$ insures that all the J -holomorphic discs are of class $\mathcal{C}^2$. There is an easy characterization of J -plurisubharmonicity in terms of some kind of Levi form (see e.g. [I-R], Corollary 1.1 page 2400), that we shall not use here, except briefly (and just for simplification) in parts IV and V. The theory for non $\mathcal{C}^2$ functions is still not completely satisfactory, although there is the interesting work of Pali [Pa].

For the basic theory of almost complex structures there are many references beyond the seminal paper [NW] ([MDF] [MDF2] [Si] in particular). Still, I shall often refer to [I-R] where special care has been taken in order to present very simple proofs. [Ro2] may also be helpful since it includes a long discussion of basic material (using the notations to be introduced later, especially of J , Q and the Nijenhuis tensor). We refer to these sources for (standard) notions or notations that will not be explained below.

Pluripolarity in almost complex structures turns out to be a somewhat delicate matter. Integrability near 0 in $\mathbf{R}^2$ of $\frac{1}{r^2 \log^2 r}$ ($r = \sqrt{x^2 + y^2}$) leads to our positive result, while non integrability of $\frac{1}{r^2 |\log r|}$ is crucial in the counterexample.

0. $\log$ and $\log |\log|$.

In this section we work on $\mathbf{C}$, with variable z .

One has $\frac{\partial^2}{\partial z \partial \bar{z}} \log |z| = 0$, on $\mathbf{C} \setminus \{0\}$.

Elementary computations give:

$$\frac{\partial^2}{\partial z \partial \bar{z}} (-\log |\log |z||) = \frac{1}{4|z|^2 \log^2 |z|} ,$$

$$\frac{\partial^2 |z|}{\partial z \partial \bar{z}} = \frac{1}{4|z|}$$

(homogeneity being obvious).

As discussed in [Ro1] the function $-\log|\log|z||$ gives an almost optimal spreading of mass for the Laplacian. Almost complex structures on $\mathbf{C}^n$ are often treated, near a point at which the structure is the standard one, as deformations of the standard complex structure. This leads to study deformations of the operator $\frac{\partial^2}{\partial z \partial \bar{z}}$ ($= \frac{1}{4}\Delta$). Corresponding to having the structure to be the standard structure at the point, one is led to consider perturbations whose terms of order 2 coincides with $\frac{\partial^2}{\partial z \partial \bar{z}}$ at 0.

Let Q be a linear differential operator defined near 0, with say smooth coefficients of the type

$$Q = \frac{\partial^2}{\partial z \partial \bar{z}} + R_2(z, D) + Q_1(z, D)$$

with $R_2(z, D) = \alpha_1(z) \frac{\partial^2}{\partial z \partial z} + \alpha_2(z) \frac{\partial^2}{\partial z \partial \bar{z}} + \alpha_3(z) \frac{\partial^2}{\partial \bar{z} \partial \bar{z}}$ and $\alpha_j(0) = 0$, $Q_1(z) = \beta_1(z) \frac{\partial}{\partial z} + \beta_2(z) \frac{\partial}{\partial \bar{z}}$, with no vanishing condition of β_j at 0.

From the estimates (for appropriate constant C)

$|\nabla(\log|\log|z||)| \leq \frac{C}{|z||\log|z||}$ and $|\nabla^2(\log|\log|z||)| \leq \frac{C}{|z|^2|\log|z||}$, it is clear that near 0:

$$Q(-|\log|\log|z||) > 0 .$$

That indicates a way that will be followed in section II.

There is no positivity of $\Delta(\log|z|)$ away from 0, so one cannot hope to keep non negativity for $Q(\log|z|)$. A corrective term is needed. Homogeneity considerations show that $C|z|^2$ cannot be enough, but we have $R_2(\log|z|)$ and $Q_1(\log|z|) = O(\frac{1}{|z|})$, $R_2(|z|)$ and $Q_1(|z|) = O(1)$. So, for $C > 0$ large enough

$$Q(\log|z| + C|z|) \geq 0 \text{ near } 0 .$$

That corresponds to the choice of the Chirka function, with logarithmic pole at a point, on almost complex manifolds, and it explains the need of having $J = J_{st}$ at the point.

In case R_2 and Q_1 vanish to second, resp. first, order at 0, then for $C > 0$ large enough one can take the function $\log|z| + C|z|^2$ (or interestingly $\log|z| + C|z - z_0|^2$ for any fixed z_0), instead of $\log|z| + C|z|$. Part IV boils down to not much more than that.

I. Preliminary Computations. Let J be an almost complex structure defined on an open set $U \subset \mathbf{C}^n$. We shall assume that J is close enough to the standard complex structure J_{st} , more precisely we assume that $J + J_{st}$ is invertible.

Then, as it is well known, the condition for J -holomorphy of a map u from say the unit disc $\mathbf{D}$ in $\mathbf{C}$ (with coordinate $\zeta = x + iy$), into U can be equivalently be given by:

(a) the ‘Cauchy-Riemann’ equation $\frac{\partial u}{\partial y} = J(u) \frac{\partial u}{\partial x}$, (where $J(u)$ is an $\mathbf{R}$ -linear map satisfying $J^2 = -\mathbf{1}$).

or

(b) an equation

$$\frac{\partial \bar{u}}{\partial \zeta} = Q(u) \frac{\partial u}{\partial \bar{\zeta}}, \quad (E1)$$

where Q is a $\mathbf{C}$ linear operator (a complex $(n \times n)$ matrix valued function) of the same smoothness as J , with $Q = 0$ whenever $J = J_{st}$. The conjugate operator $\bar{Q}$ is given by $\bar{Q} = [J + J_{st}]^{-1}[J - J_{st}]$. A long discussion of that is in [Ro2].

Let u be a J -holomorphic disc, i.e. a map from $\mathbf{D}$ into U satisfying (E1). Let λ be a real valued function defined on U , an elementary computation gives:

$$\frac{\partial^2 \lambda \circ u}{\partial \zeta \partial \bar{\zeta}} = I + II + III \quad (E2)$$

with

$$\begin{aligned} I &= \sum_{j,k \geq 1}^n \frac{\partial^2 \lambda}{\partial z_j \partial \bar{z}_k} \frac{\partial u_j}{\partial \zeta} \frac{\partial \bar{u}_k}{\partial \bar{\zeta}} + \sum_{j,k \geq 1}^n \frac{\partial^2 \lambda}{\partial z_j \partial \bar{z}_k} \frac{\partial u_j}{\partial \bar{\zeta}} \frac{\partial \bar{u}_k}{\partial \zeta} \\ II &= 2 \operatorname{Re} \sum_{j,k} \frac{\partial^2 \lambda}{\partial z_j \partial z_k} \frac{\partial u_j}{\partial \zeta} \frac{\partial u_k}{\partial \bar{\zeta}} \\ III &= 2 \operatorname{Re} \sum_j \frac{\partial \lambda}{\partial z_j} \frac{\partial^2 u_j}{\partial \zeta \partial \bar{\zeta}}. \end{aligned}$$

Standard holomorphic discs u are harmonic ($\frac{\partial^2 u}{\partial \zeta \partial \bar{\zeta}} = 0$), J -holomorphic discs of course are not, but we still have an estimate of the Laplacian in terms of the gradient. Indeed differentiation of (E1), with respect to $\bar{\zeta}$, gives:

$$\frac{\partial^2 \bar{u}}{\partial \zeta \partial \bar{\zeta}} = [Q_z(u) \cdot \frac{\partial u}{\partial \bar{\zeta}}] \frac{\partial u}{\partial \zeta} + [Q_{\bar{z}}(u) \cdot \frac{\partial \bar{u}}{\partial \bar{\zeta}}] \frac{\partial u}{\partial \zeta} + Q(u) \frac{\partial^2 u}{\partial \zeta \partial \bar{\zeta}}.$$

Since $\frac{\partial^2 \bar{u}}{\partial \zeta \partial \bar{\zeta}} = \overline{\frac{\partial^2 u}{\partial \zeta \partial \bar{\zeta}}}$, if $Q(u)$ has an operator norm $< \frac{1}{2}$, then one gets:

$$|\frac{\partial^2 u}{\partial \zeta \partial \bar{\zeta}}| \leq 2 |([Q_z(u) \cdot \frac{\partial u}{\partial \bar{\zeta}}] \frac{\partial u}{\partial \zeta} + [Q_{\bar{z}}(u) \cdot \frac{\partial \bar{u}}{\partial \bar{\zeta}}] \frac{\partial u}{\partial \zeta})|. \quad (E3)$$

Formula (E3) will be used, a first consequence that will not suffice for our purpose is that, given bounds on $|\nabla Q|$, it gives an estimate $|\frac{\partial^2 u}{\partial \zeta \partial \bar{\zeta}}| \leq C |\nabla u|^2$.

II. Pluripolarity of J -holomorphic curves.

After appropriate simple changes of variables (to straighten an embedded J -holomorphic disc and to make the almost complex structure standard along the straightened disc) the heart of the matter is the following, on which we shall now concentrate our efforts:

Proposition 1. *In $\mathbf{C}^n$, with coordinates $Z = (z_1, Z')$ ($Z' = (z_2, \dots, z_n)$), let J be a $\mathcal{C}^2$ (or $\mathcal{C}^{1,1}$) almost complex structure defined on a neighborhood of $\overline{\mathbf{D}} \times \{0\}$ (the set $|z_1| \leq 1$, $z_2 = \dots = z_n = 0$). Assume that $J = J_{st}$ along $\mathbf{C} \times \{0\}$. Then for $K > 0$ large enough the function*

$$\Lambda(Z) = -\log |\log |Z'||| + K|z_1|^2$$

is J -plurisubharmonic on a neighborhood of $\overline{\mathbf{D}} \times \{0\}$.

Proof. Set $LL(Z') = -\log |\log |Z'|||$. The starting point is the estimate for the standard complex Hessian of the function $LL(Z')$. See II.1 in [Ro1]

$$4 \sum_{j,k \geq 2} \frac{\partial^2 LL(Z')}{\partial z_j \partial \bar{z}_k} t_j \bar{t}_k \geq \frac{\sum_{j \geq 2} |t_j|^2}{|Z'|^2 \log^2 |Z'|} .$$

Let u be a J -holomorphic disc (with values near $\overline{\mathbf{D}} \times \{0\}$), we want to show that $\frac{\partial^2}{\partial \zeta \partial \bar{\zeta}} \Lambda \circ u(0) \geq 0$.

(1) We first estimate $\frac{\partial^2}{\partial \zeta \partial \bar{\zeta}} LL \circ u(0)$, by taking $\lambda(Z) = LL(Z')$ in (E2). We shall estimate separately the terms I, II, and III, of course in the summations only j and $k \geq 2$ need to be considered. Various constants will be denoted by the same letter C in the proof.

Set $u(0) = (z_1^0, Z'_0)$, $\tau = (\sum_{j=2}^n |\frac{\partial u_j}{\partial \zeta}(0)|^2)^{\frac{1}{2}}$, and $\epsilon = |\frac{\partial u_1}{\partial \zeta}(0)|$. For the $\bar{\zeta}$ derivatives, for some appropriate constant C , one has the estimate $|\frac{\partial u_j}{\partial \bar{\zeta}}(0)| \leq C|Z'_0|(\epsilon + \tau)$, according to (E1) and since Q is of class $\mathcal{C}^1$ and vanishes for $Z' = 0$. It is an essential difficulty that (E1) does not give a control of the $\bar{\zeta}$ derivatives of the u_j 's for $j \geq 2$ only in terms of the ζ derivatives of these functions alone, $\frac{\partial u_1}{\partial \zeta}$ comes in the estimate. However, only the partial derivatives $\frac{\partial u_j}{\partial \zeta}(0)$ for $j \geq 2$ contribute to the initial crucial positivity in the proof.

Estimate of I. We have

$$I \geq \frac{\tau^2}{4|Z'_0|^2 \log^2(|Z'_0|)} .$$

Estimate of II. For an (other) appropriate constants C , $|\nabla^2 LL(Z')| \leq \frac{C}{|Z'|^2 |\log |Z'||}$. So, for II we get (for some other appropriate constant C):

$$|II| \leq \frac{C}{|Z'_0| |\log |Z'_0||} (\epsilon + \tau) \tau .$$

Estimate of III. One first has an estimate $|\nabla LL(Z')| \leq \frac{C}{|Z'| |\log |Z'||}$. More delicate is the estimate of $\frac{\partial^2 u}{\partial \zeta \partial \bar{\zeta}}(0)$. Here we need to analyze both terms on the right hand side of (E3). Using that $Q = 0$ if $Z' = 0$, that gives $|\frac{\partial u}{\partial \zeta}(0)| \leq C|Z'_0| \frac{\partial u}{\partial \zeta}(0) = C|Z'_0|(\epsilon + \tau)$, one gets

$$[Q_z(u) \cdot \frac{\partial u}{\partial \bar{\zeta}}] \frac{\partial u}{\partial \zeta}(0) \leq C|Z'_0|(\epsilon^2 + \tau^2) .$$

Next we evaluate (for $\zeta = 0$) $Q_{\bar{z}}(u) \cdot \frac{\partial u}{\partial \bar{\zeta}} = a + b$, where $a = \frac{\partial u_1}{\partial \zeta} \frac{\partial Q}{\partial \bar{z}_1}$ and $b = \sum_{j=2}^n \frac{\partial u_j}{\partial \zeta} \frac{\partial Q}{\partial \bar{z}_j}$. Since $Q \equiv 0$ for $Z' = 0$, one has $|\frac{\partial Q}{\partial \bar{z}_1}(Z)| \leq C|Z'|$, and this is where we need $\mathcal{C}^2$ regularity of Q (or at least $\mathcal{C}^{1,1}$ regularity). Then one gets $|a| \leq C|Z'_0|\epsilon$. For the b term, one sees that $|b| \leq C\tau$. Therefore $|[Q_{\bar{z}}(u) \cdot \frac{\partial u}{\partial \bar{\zeta}}] \frac{\partial u}{\partial \zeta}(0)| \leq C(|Z'_0|\epsilon + \tau)(\epsilon + \tau)$. Consequently

$$|\frac{\partial^2 u}{\partial \zeta \partial \bar{\zeta}}(0)| \leq C|Z'_0|(\epsilon^2 + \tau^2) + C\tau(\epsilon + \tau) .$$

Finally

$$|III| \leq C(\frac{\epsilon^2}{|\log |Z'_0||} + \frac{\tau^2}{|Z'_0| |\log |Z'_0||} + \frac{\epsilon\tau}{|Z'_0| |\log |Z'_0||}) .$$

Putting together all the estimates obtained so far, we get (for new constants C):

$$\begin{aligned} \frac{\partial^2}{\partial \zeta \partial \bar{\zeta}} LL \circ u (0) &\geq \frac{\tau^2}{4|Z'_0|^2 \log^2 |Z'_0|} \\ &- \frac{C\tau^2}{|Z'_0| |\log |Z'_0||} - \frac{C\epsilon\tau}{|Z'_0| |\log |Z'_0||} - \frac{C\epsilon^2}{|\log |Z'_0||} . \end{aligned}$$

On the right hand side, the first term gives positivity, and the worst term is $-\frac{C\epsilon\tau}{|Z'_0| |\log |Z'_0||}$.

Remark: Before finishing the proof it is good to point out that if $\epsilon = 0$, the right hand side is clearly non negative provided $|Z'_0|$ is small enough, i.e. $u(0)$ is close to $\mathbf{C} \times \{0\}$. In fact the estimates above prove that in $\mathbf{C}^N$ equipped with an almost structure J of class $\mathcal{C}^1$, with $J(0) = J_{st}$, the function $-\log |\log |Z||$ is J -plurisubharmonic near 0. No additional corrective term is needed. The proof is just a matter of dropping all the terms with index $j = 1$ above. In particular the term a in the estimate of III above need not be considered and this is why $\mathcal{C}^1$ smoothness would be enough. Also, note that at any rate J -purisubharmonicity of $-\log |\log |Z||$ is, for $\mathcal{C}^2$ data, a completely trivial consequence of the Proposition, by consideration of $\mathbf{C}^N \times \mathbf{C}_{st}$.

(2) We now estimate $\frac{\partial^2}{\partial \zeta \partial \bar{\zeta}} |u_1|^2$. So this time we apply (E2) to the function $\lambda(Z) = |Z_1|^2$. With the notation as in (1), one gets immediately

$$I \geq \epsilon^2 \quad , \quad II = 0 .$$

For III, $\frac{\partial \lambda}{\partial z_1} = \bar{z}_1$, and $\frac{\partial \lambda}{\partial z_j} = 0$ if $j \neq 1$, but we need to estimate $\frac{\partial^2 u_1}{\partial \zeta \partial \bar{\zeta}}$ as in (1). This gives us

$$|III| \leq C(|Z'_0|\epsilon^2 + \tau^2 + \epsilon\tau) \quad .$$

(3) According to (1) and (2) $\Lambda = LL(Z') + K|Z_1|^2$ will be J -plurisubharmonic near $\bar{\mathbf{D}} \times \{0\}$ if for $|Z'_0|$ small enough, and for any ϵ and $\tau \geq 0$,

$$\begin{aligned} & \frac{\tau^2}{4|Z'_0|^2 \log^2 |Z'_0|} + K\epsilon^2 \\ & - \frac{C\tau^2}{|Z'_0| \log |Z'_0|} - \frac{C\epsilon\tau}{|Z'_0| \log |Z'_0|} - \frac{C\epsilon^2}{|\log |Z'_0||} - CK|Z'_0|\epsilon^2 - CK\tau^2 - CK\epsilon\tau \geq 0 \quad . \end{aligned}$$

By taking $K > 4C^2$, one has

$$\frac{\tau^2}{4|Z'_0|^2 \log^2 |Z'_0|} + K\epsilon^2 - \frac{C\epsilon\tau}{|Z'_0| \log |Z'_0|} \geq \frac{1}{8} \left(\frac{\tau^2}{|Z'_0|^2 \log^2 |Z'_0|} + K\epsilon^2 \right) \quad .$$

Once K has been fixed, the other terms are immediately absorbed when $|Z'_0|$ is small enough. This ends the proof of the Proposition.

Remark: To get a logarithmic singularity, one would be tempted to add $c \log |Z'|$ to Λ , for $c > 0$ small enough. Then, coming from term II in E2, and from estimates of derivatives, a term arises with $\frac{1}{|Z'_0|}$ instead of $\frac{1}{|Z'_0| |\log |Z'_0||}$. As shown in Part III, this term cannot be controlled.

III. An Example.

On $\mathbf{C}^2$, we consider the almost complex structure J defined by taking

$$Q = \begin{pmatrix} 0 & 0 \\ z_2 & 0 \end{pmatrix} \quad .$$

So the equations for J -holomorphic discs $\zeta \mapsto u(\zeta) = (u_1(\zeta), u_2(\zeta))$ are: u_1 must be holomorphic, and $\frac{\partial \bar{u}_2}{\partial \bar{\zeta}} = u_2 \frac{\partial u_1}{\partial \zeta}$.

It then follows that $\frac{\partial u_2}{\partial \zeta} = \bar{u}_2 \frac{\partial \bar{u}_1}{\partial \bar{\zeta}}$, and $\frac{\partial^2 u_2}{\partial \zeta \partial \bar{\zeta}} = u_2 \left| \frac{\partial u_1}{\partial \zeta} \right|^2$.

The fact that $Q_z \neq 0$ is important, this is related to the non vanishing of the Nijenhuis tensor. See detailed discussion in [Ro2] (Proposition 6), and IV below. A direct description of J is given at the end of Part III.

Note that $J = J_{st}$ along $\mathbf{C} \times \{0\}$, that is therefore J -holomorphic.

Proposition 2. *There is no J -plurisubharmonic function λ , defined near 0 and not identically $-\infty$ near 0, such that $\lambda(z_1, z_2) \leq \log |z_2|$.*

Recall that, for plurisubharmonic functions, we restrict our attention to functions that are $\mathcal{C}^2$ off their $-\infty$ set. I have not tried to extend Proposition 2 to the case of merely continuous plurisubharmonic functions. The proof will be by contradiction, so we assume that there is a function λ with the desired properties.

We rewrite equation (E3), that simplifies, and with a different arrangement of terms, Let $u = (u_1, u_2)$ be a J -holomorphic map from a neighborhood of 0 in $\mathbf{C}$ into some fixed neighborhood of 0 in $\mathbf{C}^2, J$).

$$\text{Set } u(0) = (z_1, z_2) \text{ . Assume } \frac{\partial u_1}{\partial \zeta}(0) = 1 \text{ , set } \frac{\partial u_2}{\partial \zeta}(0) = t \text{ .}$$

Note that except if $\frac{\partial u_1}{\partial \zeta}(0) = 0$, $\frac{\partial u_1}{\partial \zeta}(0) = 1$ can be achieved by simple rescaling. By the equations for J -holomorphy: $\frac{\partial u_1}{\partial \bar{\zeta}} = 0$, and $\frac{\partial u_2}{\partial \bar{\zeta}}(0) = \bar{z}_2$. One gets

$$\frac{\partial^2}{\partial \zeta \partial \bar{\zeta}}(\lambda \circ u)(0) = A + B + C_1 + C_2 \quad (E4)$$

with:

$$\begin{aligned} A &= 2 \operatorname{Re}[(\frac{\partial^2 \lambda}{\partial \bar{z}_1 \partial z_2} + \frac{\partial^2 \lambda}{\partial z_2^2} \bar{z}_2)t] \text{ ,} \\ B &= \frac{\partial^2 \lambda}{\partial z_2 \partial \bar{z}_2} (|t|^2 + |z_2|^2) \text{ ,} \\ C_1 &= 2 \operatorname{Re}[(\frac{\partial^2 \lambda}{\partial z_1 \partial z_2} \bar{z}_2) + (\frac{\partial \lambda}{\partial z_2} z_2)] \text{ ,} \\ C_2 &= \frac{\partial^2 \lambda}{\partial z_1 \partial \bar{z}_1} \text{ .} \end{aligned}$$

The strategy for proving the Proposition will be to find $K_1 > 0$ such that for $z_1 = 0$ and some z_2 's, with $|z_2|$ arbitrarily small, one can choose t such that $A + B + C_1 + C_2 \leq K_1 \log |z_2| < 0$. Negativity will come from the A term, more precisely from $\frac{\partial^2 \lambda}{\partial z_2^2} \bar{z}_2$, coming from II which was previously treated as an error term. In our example, one can avoid cancellation by $\frac{\partial^2 \lambda}{\partial \bar{z}_1 \partial z_2}$ by appropriate choice of the argument of z_2 . The other terms will be 'error terms'.

It happens that a good choice of t will be with $|t| = K_2 |z_2| |\log |z_2||$, for some appropriate value of K_2 .

Preliminary Remarks.

For any fixed $z_1 \in \mathbf{C}$, $\zeta \mapsto (z_1, \zeta)$ is a J -holomorphic map. Hence $z_2 \mapsto \lambda(z_1, z_2)$ is a subharmonic function. Because of the Logarithmic singularity we must have

$$\lambda(z_1, z_2) = a(z_1) \log |z_2| + \mu(z_1, z_2) \text{ ,} \quad (E4)$$

where $a(z_1) \geq 1$ (the Lelong number) and the measure ν_{z_1} , on $\mathbf{C}$ (near 0), defined by $\nu_{z_1} = \frac{\partial^2}{\partial z_2 \partial \bar{z}_2}(\mu(z_1, \cdot))$ has no point mass at $z_2 = 0$. So, $\mu(z_1, z_2)$ can be thought as a term of lesser singularity along $z_2 = 0$.

1. We shall first prove the non existence of λ if one imposes some conditions that seem very natural and reasonable, in view of the above discussion.

The hypotheses that we now make on the term μ , 'less singular' than $\log|z_2|$, are:

For z_1 small, and as $z_2 \rightarrow 0$:

$$z_2^2 \frac{\partial^2 \mu}{\partial z_2^2} \rightarrow 0 \quad , \quad z_2 \frac{\partial \mu}{\partial z_2} \rightarrow 0 \quad . \quad (\mathbf{H})$$

Since μ is real valued the second condition implies that $\frac{\mu}{\log|z_2|} \rightarrow 0$, but we won't need this remark.

Since we clearly have invariance under z_1 translations, we can partially smooth λ (and thus a and μ) by convolution in the z_1 variable. Then $a(\cdot)$ is smooth and for fixed z_2 , a sup norm estimate of λ or μ or of some derivative of these functions, results in a similar sup norm estimate for functions obtained by further differentiations in the z_1 direction. We can therefore assume that additionally to $(\mathbf{H})$, we also have, for some constant $M > 0$:

$$|\frac{\partial^2 \lambda}{\partial z_1 \partial \bar{z}_1}| \leq M |\log|z_2|| \quad , \quad |\frac{\partial^2 \lambda}{\partial z_1 \partial z_2}| \leq \frac{M}{|z_2|} \quad , \quad z_2 \frac{\partial^2 \mu}{\partial \bar{z}_1 \partial z_2} \rightarrow 0, \text{ as } z_2 \rightarrow 0 \quad . \quad (\mathbf{H}^+)$$

Proof of Proposition assuming $(\mathbf{H})$ (and thus $(\mathbf{H}^+)$).

We first evaluate $A + C_1 + C_2$. We have

$$\begin{aligned} \frac{\partial^2 \lambda}{\partial \bar{z}_1 \partial z_2} &= \frac{1}{2} \frac{\partial a}{\partial \bar{z}_1} \frac{1}{z_2} + \frac{\partial^2 \mu}{\partial \bar{z}_1 \partial z_2} = \frac{1}{2} \frac{\partial a}{\partial \bar{z}_1} \frac{1}{z_2} + o\left(\frac{1}{|z_2|}\right) , \\ \frac{\partial^2 \lambda}{\partial z_2^2} &= \frac{-a(z_1)}{2z_2^2} + o\left(\frac{1}{|z_2|^2}\right) . \end{aligned}$$

Consequently

$$\frac{\partial^2 \lambda}{\partial \bar{z}_1 \partial z_2} + \frac{\partial^2 \lambda}{\partial z_2^2} \bar{z}_2 = \frac{1}{2} \left(\frac{\partial a}{\partial \bar{z}_1} - a(z_1) \frac{\bar{z}_2}{z_2} \right) \frac{1}{z_2} + o\left(\frac{1}{|z_2|}\right) .$$

For any fixed z_1 , and we take $z_1 = 0$, there exists an interval $\mathcal{I}$ in $\mathbf{R}/2\pi\mathbf{Z}$, of length $|\mathcal{I}| \geq \frac{\pi}{4}$, such that if $z_2 = re^{i\theta}$ with $\theta \in \mathcal{I}$, (avoiding cancellation of $-a$ by $\frac{\partial a}{\partial \bar{z}_1}$) one has

$$|\frac{\partial^2 \lambda}{\partial \bar{z}_1 \partial z_2} + \frac{\partial^2 \lambda}{\partial z_2^2} \bar{z}_2| \geq \frac{1}{3|z_2|} \quad ,$$

if $|z_2|$ is small enough.

Under our hypotheses, as z_2 approaches 0, the C_1 term stays bounded and for the C_2 term one gets a bound $|C_2| \leq M|\log|z_2||$. Therefore if we take K large enough, for any $z_2 = |z_2|e^{i\theta}$, with $\theta \in \mathcal{I}$, we can choose t with $|t| = K|z_2||\log|z_2||$, with the argument chosen so that

$$A + C_1 + C_2 \leq -\frac{K}{4} |\log|z_2||.$$

We finally need to control the B term. Ideally we would like $(|z_2|^2 \log|z_2|) \frac{\partial^2 \lambda}{\partial z_2 \partial \bar{z}_2} \rightarrow 0$. That would make the B term negligible. Note that this is asking more than the natural requirement suggested by ‘homogeneity’ consideration, that would be $(|z_2|^2) \frac{\partial^2 \lambda}{\partial z_2 \partial \bar{z}_2} \rightarrow 0$, since $\frac{\partial^2 \lambda}{\partial z_2 \partial \bar{z}_2} = \frac{\partial^2 \mu}{\partial z_2 \partial \bar{z}_2}$.

Since, as pointed out earlier, λ is a subharmonic function of z_2 (for z_1 fixed), its Laplacian is a bounded positive measure. For fixed z_1 , $\frac{\partial^2 \lambda}{\partial z_2 \partial \bar{z}_2}$ is (near $z_2 = 0$) the sum of a point mass at 0 and of an integrable function. So, given any $\epsilon > 0$ the inequality $|\frac{\partial \lambda}{\partial z_2 \partial \bar{z}_2}| > \frac{\epsilon}{|z_2|^2 |\log|z_2||}$ cannot be satisfied near 0 by all z_2 in a given sector. Then, at any point z_2 where the reverse inequality $|\frac{\partial \lambda}{\partial z_2 \partial \bar{z}_2}| \leq \frac{\epsilon}{|z_2|^2 |\log|z_2||}$ holds, for t as above one has

$$|B| = |\frac{\partial^2 \lambda}{\partial z_2 \partial \bar{z}_2} (|t|^2 + |z_2|^2)| \leq \frac{\epsilon}{|z_2|^2 |\log|z_2||} K^2 |z_2|^2 \log^2 |z_2| + \frac{\epsilon}{|\log|z_2||} \leq \epsilon K^2 |\log|z_2|| + 1.$$

By taking ϵ so that $\epsilon K^2 < \frac{K}{8}$, one gets $A + B + C_1 + C_2 \leq -\frac{K}{8} |\log|z_2|| + 1 < 0$, ($|z_2|$ small). Therefore λ is not J -plurisubharmonic since its restriction to any J -holomorphic disc satisfying $u(0) = (0, z_2)$, $\frac{\partial u_1}{\partial \zeta}(0) = 1$ and $\frac{\partial u_2}{\partial \zeta}(0) = t$, will not be subharmonic.

2. We now prove Proposition 3 in full generality. Our first task is to get that $z_2 \frac{\partial \mu}{\partial z_2}$ tends to 0, uniformly with respect to z_1 , as z_2 approaches zero along some ‘fat’ set.

A subset $E \subset \mathbf{C}$ will be said to be *fat* if 0 is a point of Lebesgue density of E , i.e.

$$\frac{|E \cap \{|z| < r\}|}{\pi r^2} \rightarrow 1 \quad \text{as } r \rightarrow 0.$$

Assume that there is a function λ_1 with the properties mentioned in the statement of the Proposition, defined for $|z_1| \leq 4R$, $|z_2| \leq 3R$. For $0 < |z_2| < 2R$ (so $z_2 \neq 0$), set

$$\nu(z_2) = \int_{|z_1| < 3R} \frac{\partial^2 \lambda_1}{\partial z_2 \partial \bar{z}_2}(z_1, z_2) dm(z_1)$$

where dm denotes Lebesgue measure. By subharmonicity of λ_1 in the variable z_2 , ν is an integrable function. Let λ be a function obtained by partially smoothing λ_1 by

convolution in the z_1 variable, with a non non negative function of z_1 , of small support. J -plurisubharmonicity is preserved since J is invariant under z_1 translations, and we replace the original function λ_1 by λ . Then there is $C > 0$ such that for all $|z_1| < 2R$ we have

$$0 \leq \frac{\partial^2 \lambda}{\partial z_2 \partial \bar{z}_2}(z_1, z_2) \leq C\nu(z_2) .$$

Let μ be the function defined in (E4). Let $\chi(z_2)$ be the characteristic function of the set $\{|z_2| < 2R\}$. Then for each fixed z_1 , on the set $|z_2| < R$, μ is the sum of a smooth function (with uniform bounds) and of $(\chi \frac{\partial^2 \mu}{\partial z_2 \partial \bar{z}_2}) * \frac{2 \log |z_2|}{\pi}$ (convolution, in the z_2 variable, of the Laplacian with the Newtonian potential). And $\frac{\partial^2 \mu}{\partial z_2 \partial \bar{z}_2}$ is simply the restriction to $z_2 \neq 0$ of $\frac{\partial \lambda}{\partial z_2 \partial \bar{z}_2}$ (the point mass at $\{0\}$ being thus dropped). So, modulo smooth function $\frac{\partial \mu}{\partial z_2} = (\chi \frac{\partial^2 \mu}{\partial z_2 \partial \bar{z}_2}) * \frac{1}{\pi z_2}$. We therefore get that for every $|z_1| < 2R$, for $|z_2| < R$

$$|\frac{\partial \mu}{\partial z_2}| \leq C \frac{1}{|z_2|} * \nu + C$$

where the convolution takes place in the z_2 variable. It then follows from Lemma A1 in the appendix that there is a fat set E such that:

$$\text{Sup}_{|z_1| \leq R} |z_2 \frac{\partial \mu}{\partial z_2}(z_1, z_2)| \rightarrow 0 , \text{ as } z_2 \rightarrow 0 , z_2 \in E .$$

So we have a replacement of the second hypothesis in **H**, if we restrict z_2 to E .

It is essential that for $z_2 \in E$ we had estimates, uniformly for $|z_1| < R$. As done in **1** we can replace λ by a function obtained by a second (probably un-needed) smoothing by convolution in the z_1 variable and if one restricts to $z_2 \in E$, hypotheses **H**⁺ are satisfied.

We finally need a replacement for the first hypothesis in **H**. This is a more subtle matter since one has to consider now a singular integral.

$$\frac{\partial^2 \mu}{\partial z_2^2} = \nu * \frac{-1}{\pi z_2^2} + g,$$

where g is a function bounded near 0.

For the end of the proof, contrary to above, we have to fix z_1 . Take $z_1 = 0$. Lemma A2 in the Appendix shows that there exists a fat set E' ($\subset E$) on which additionally to the above we have $z_2^2 \frac{\partial^2 \mu}{\partial z_2^2} \rightarrow 0$ as $z_2 \rightarrow 0$, $z_2 \in E'$.

Lemma A3 asserts that for any $\epsilon > 0$ the inequality $\frac{\partial^2 \lambda}{\partial z_2 \partial \bar{z}_2} > \frac{\epsilon}{|z|^2 \log |z|}$ cannot hold for all z_2 near 0 $z_2 \neq 0$, and $z_2 \in E'$. From here the proof is identical to the proof in **1**, with the only difference that one has to take $z_2 \in E'$.

3. Tensor J . The almost complex structure of the example on $\mathbf{C}^2$ has been given by the $\mathbf{C}$ -linear operator Q that characterizes J -holomorphy:

$$\frac{\partial \bar{u}}{\partial \zeta} = Q(u) \frac{\partial u}{\partial \zeta} .$$

We now wish to write the corresponding $\mathbf{R}$ linear operator J defining the almost complex structure on $\mathbf{R}^4$, on which coordinates will be (x_1, y_1, x_2, y_2) (identified with $\mathbf{C}^2$, by setting $z_j = x_j + iy_j$).

There is a formula: $J = J_{st}[\mathbf{1} - \bar{Q}][\mathbf{1} + \bar{Q}]^{-1}$ (see 2.1 in [Ro2]) that one could use. We do a direct computation instead. We observe that $\zeta = (x + iy) \mapsto (z_1 + \zeta, z_2 + \bar{z}_2 \bar{\zeta})$ satisfies the equation for J -holomorphicity at $\zeta = 0$. In terms of J the equation is $\frac{\partial u}{\partial y} = J(u) \frac{\partial u}{\partial x}$. Therefore

$$[J(z_1, z_2)](1, \bar{z}_2) = (i, -i\bar{z}_2) .$$

Since $J^2 = -\mathbf{1}$, $[J(z_1, z_2)](i, -i\bar{z}_2) = (-1, -\bar{z}_2)$.

Since $\zeta \mapsto (z_1, z_2 + \zeta)$ is J -holomorphic, one also has

$$[J(z_1, z_2)](0, \xi) = (0, i\xi) , \text{ for any } \xi \in \mathbf{C} .$$

One immediately gets

$$[J(z_1, z_2)](1, 0) = (i, -2i\bar{z}_2) , \quad [J(z_1, z_2)](i, 0) = (-1, -2\bar{z}_2) ,$$

$$[J(z_1, z_2)](0, 1) = (0, i) , \quad [J(z_1, z_2)](0, i) = (-1, 0) .$$

With real $\mathbf{R}^4$ notations at point (x_1, y_1, x_2, y_2) , J is given by the matrix

$$J = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ -2y_2 & -2x_2 & 0 & -1 \\ -2x_2 & 2y_2 & 1 & 0 \end{pmatrix} .$$

IV. Logarithmic singularities.

There is no surprise that if we assume better matching with the standard complex structure along an embedded J -holomorphic curve, there will be a J -plurisubharmonic function with logarithmic singularity along that curve. Better matching with the standard complex structure is given by vanishing of the Nijenhuis tensor.

On an almost complex manifold M a $(0, 1)$ vector field is a complexified tangent vector field $\bar{L}$ such that $\bar{L} = X + iJX$ for some real tangent vector field. The vanishing of the Nijenhuis tensor at a point $p \in M$ is equivalent to the fact that if $\bar{L}_1$ and $\bar{L}_2$ are $(0, 1)$ vector fields, their Lie bracket is of type $(0, 1)$ at p (i.e. $[\bar{L}_1, \bar{L}_2](p) = X_p + iJ(p)X_p$, for

some real tangent vector X_p to M at p). Although smoothness hypotheses are made precise below, we did not try to improve them, e.g. by using the Whitney extension Theorem.

Lemma. *Let J be an almost complex structure of class $\mathcal{C}^4$ defined on a neighborhood of $\overline{\mathbf{D}} \times \{0\}$ in $\mathbf{C}^n$. Assume that $J = J_{st}$ along $\mathbf{C} \times \{0\}$. The following are equivalent:*

- (a) *The Nijenhuis tensor N_J vanishes along $\mathbf{D} \times \{0\}$*
- (b) *There exists a $\mathcal{C}^3$ local change of coordinates given by:*

$$Z_r(z) = z_r + \sum_{k,l \geq 2} a_{k,l}^r(z_1) z_k \bar{z}_l + \sum_{k,l \geq 2} b_{k,l}^r(z_1) \bar{z}_k \bar{z}_l ,$$

such that for any $\mathcal{C}^2$ $(0,1)$ vector field $\bar{L}$

$$\bar{L}(Z_j)(z_1, z_2, \dots, z_n) = O\left(\sum_{j \geq 2} |z_j|^2\right) ,$$

in a neighborhood of $\mathbf{D} \times \{0\}$.

The proof is identical to the proof of Proposition 4 in [Ro2], with z_1 as a parameter. In the new coordinates the almost complex structure is of class $\mathcal{C}^2$ at least. We shall prove that (a) implies (b). The converse is easy since ∇Q , hence $\nabla J = 0$ in the new coordinates, and evaluating the Nijenhuis tensor requires only one derivative of J .

Proof: By simple linear algebra there is a basis of $(0,1)$ vector fields $\bar{L}_j$ ($j = 1, \dots, n$) such that

$$\bar{L}_j = \frac{\partial}{\partial \bar{z}_j} + \sum_q \alpha_{j,q}(z) \frac{\partial}{\partial z_q} ,$$

where the functions $\alpha_{j,q}$ vanish along $\mathbf{D} \times \{0\}$. (a) reduces to the simple form $[\bar{L}_j, \bar{L}_k] = 0$ along $\mathbf{D} \times \{0\}$, giving us

$$\frac{\partial \alpha_{j,q}}{\partial \bar{z}_k}(z_1, 0, \dots) = \frac{\partial \alpha_{k,q}}{\partial \bar{z}_j}(z_1, 0, \dots) .$$

We have

$$\bar{L}_j(Z_r) = \alpha_{j,r}(z) + \sum_{k \geq 2} a_{k,j}^r(z_1) z_k + \sum_{k \geq 2} (b_{k,j}^r(z_1) + b_{j,k}^r(z_1)) \bar{z}_k + O\left(\sum_{j \geq 2} |z_j|^2\right) .$$

We then choose $a_{k,j}^r(z_1) = -\frac{\partial \alpha_{j,r}}{\partial z_k}(z_1, 0, \dots)$ and $b_{k,j}^r(z_1)$ such that $b_{k,j}^r(z_1) + b_{j,k}^r(z_1) = -\frac{\partial \alpha_{j,r}}{\partial \bar{z}_k}(z_1, 0, \dots)$. This is possible since $\frac{\partial \alpha_{j,q}}{\partial \bar{z}_k}(z_1, 0, \dots) = \frac{\partial \alpha_{k,q}}{\partial \bar{z}_j}(z_1, 0, \dots)$.

In the coordinates (Z_j) of the Lemma, J coincides to J_{st} to second order along $\mathbf{D} \times \{0\}$. So, along $\mathbf{D} \times \{0\}$, not only $Q = 0$ but also $\nabla Q = 0$. Recall that $Q_{\bar{z}} = 0$ can always be achieved according to Sukhov and Tumanov (see [D-S] Lemma 3.2, sections 2.2 in [S-T1])

[S-T2], and [Ro2] in particular Proposition 6). It is for getting $Q_z = 0$ that one needs the vanishing of the Nijenhuis tensor.

Proposition 3. *In $\mathbf{C}^n$, with coordinates $Z = (z_1, Z')$ ($Z' = (z_2, \dots, z_n)$), let J be a $\mathcal{C}^4$ almost complex structure defined on a neighborhood of $\overline{\mathbf{D}} \times \{0\}$ (the set $|z_1| \leq 1, z_2 = \dots = z_n = 0$). Assume that along $\mathbf{C} \times \{0\}$, $J = J_{st}$ and the Nijenhuis tensor N_J vanishes. Then for $K > 0$ large enough the function*

$$\Lambda(Z) = \log |Z'| + K|Z|^2$$

is J -plurisubharmonic on a neighborhood of $\overline{\mathbf{D}} \times \{0\}$.

Proof. The proof is similar to the proof in part **II**, and simpler. Due to the second order vanishing of Q along $\mathbf{C} \times \{0\}$. Quickly said: $\log |Z'| \circ u$ (that would be subharmonic if u were holomorphic) has a Laplacian bounded from below, if u is J -holomorphic, and $|\nabla u| \leq 1$. Then $|Z|^2$ brings enough positivity.

We adopt the same notations as in the Proof of Proposition 3. We estimate $\frac{\partial^2(\log |z'| \circ u)}{\partial \zeta \partial \bar{\zeta}}(0)$ by taking $\lambda = \log |Z'|$ in (E2). We have $I \geq 0$ since $\log |Z|$ is J_{st} -plurisubharmonic.

For II $|\frac{\partial u_k}{\partial \bar{\zeta}}|(0) \leq C|Z'_0|^2(\epsilon + \tau)$ (with now $|Z'_0|^2$ instead of $|Z'_0|$), but now $|\nabla^2 \lambda| \leq \frac{C}{|Z'_0|^2}$, so (with a new constant C) $II \leq C(\epsilon + \tau)\tau$.

For III, there is a gain in estimating the b term in the proof of Part **II**, we now have $\frac{\partial Q}{\partial \bar{z}_j} = 0$ for $Z' = 0$, not only for $j = 1$. This results in having $|\frac{\partial^2 u}{\partial \zeta \partial \bar{\zeta}}(0)| \leq C|Z'_0|(\epsilon^2 + \tau^2)$. Thus $|III| \leq C(\epsilon^2 + \tau^2)$ and finally

$$\frac{\partial^2 \log(|Z'| \circ u)}{\partial \zeta \partial \bar{\zeta}}(0) \geq -C(\epsilon^2 + \tau^2).$$

If Z'_0 is close to 0, one has $\frac{\partial^2(|Z'|^2 \circ u)}{\partial \zeta \partial \bar{\zeta}}(0) \geq M(\epsilon^2 + \tau^2)$ (M a positive constant). This can be obtained by the same considerations as above or more simply by using that along $\mathbf{C} \times \{0\}$ $dd^c_j = dd^c$ and the simple formula

$$\Delta(\lambda \circ u)(\zeta) = [dd^c_j \lambda]_{u(\zeta)}\left(\frac{\partial u}{\partial x}, [J(u)]\frac{\partial u}{\partial x}\right),$$

see e.g. Lemma 1.2 in [I-R].

If K is large enough, one therefore gets $\frac{\partial^2(\Lambda \circ u)}{\partial \zeta \partial \bar{\zeta}}(0) \geq 0$. Proposition 3 is thus proved.

V. Non-pluripolarity of generic submanifolds.

It is often the case that an argument using Jensen measures can be avoided and can be replaced by an elementary reasoning. However (at least for people with some training in function algebras) Jensen measures give an immediate insight. So we start by discussing

Jensen measures although we shall say later how elementary reasoning suffices (so **1** can be skipped).

1) Jensen Measures. No originality can be claimed for the following:

Lemma 2. *Let K be a compact set in an almost complex manifold M , and let $p \in M$. Assume that for every J -plurisubharmonic function λ , $\lambda(p) \leq \text{Sup}_K \lambda$. Then there exists a probability measure μ on K such that for every J -plurisubharmonic function λ , $\lambda(p) \leq \int_K \lambda d\mu$.*

Such a measure will be called a Jensen measure for p on K . The proof is the standard proof of the existence of Jensen measures in the theory of uniform algebras (see e.g. [Ga], or [St] page 11).

Proof. Since from the beginning we have restricted our attention to J -plurisubharmonic functions λ that are continuous off their $-\infty$ set, we can restrict our attention to those that are continuous and that take values in $(-\infty, +\infty)$, approximating λ by $\chi_N \circ \lambda$, where χ_N is a convex increasing function defined on $\mathbf{R}$ that satisfies $\chi_N(t) = -N$ if $t \leq -N - 1$ and $\chi_N(t) = t$ if $t \geq -N + 1$, with $N \rightarrow +\infty$.

Let Γ be the open convex cone in the space of real valued continuous functions on K that consists of the functions φ for which there is a J -plurisubharmonic function λ defined on M , such that $\lambda(p) > 0$ and $\lambda \leq \varphi$ on K . Then $0 \notin \Gamma$. By the Hahn-Banach Theorem, there exists a linear functional ϕ on the space of continuous functions on K , that is positive on Γ . All positive functions on K belong to Γ (by taking constant functions for λ), so ϕ is a positive functional. After multiplication by appropriate constant, ϕ is given by a probability measure μ . If $\alpha \in \mathbf{R}$ and λ is a J -plurisubharmonic function such that $\alpha < \lambda(p)$, then $(\lambda - \alpha) \in \Gamma$, so $\int_K (\lambda - \alpha) d\mu > 0$. Hence $\alpha < \int_K \lambda d\mu$ since $\int_K d\mu = 1$. Q.E.D.

2) Dimension 2. First, we show that if λ is a (standard) plurisubharmonic function defined near 0 in $\mathbf{C}^2$ that is $-\infty$ on $\mathbf{R}^2$ near 0, then λ is identically $-\infty$ on $\mathbf{C}^2$ near 0. There are of course many immediate arguments for showing that. The one here is not the simplest one (by far) but the advantage is that it extends immediately to the almost complex setting, by treating the almost complex case, just as a small perturbation of the complex case, as it is usual to do, with no delicate analysis needed this time.

For $r > 0$ small enough let $U = \{(z_1, z_2), \text{Im } z_1 > 0, |\text{Im } z_2| \leq (\text{Im } z_1)^2, |z_1|^2 + |z_2|^2 < r^2\}$. The boundary of U consists of 3 parts A, B and C : with $A \subset \mathbf{R}^2$, B a subset of the sphere of radius r , and $C \subset \{\text{Im } z_1 > 0, |\text{Im } z_2| = (\text{Im } z_1)^2\}$. C is strictly pseudo concave (tube domain with concave boundary), so for any plurisubharmonic function λ defined on the closure of U , $\text{Sup}_U \lambda = \text{Sup}_{A \cup B} \lambda$. For $p \in U$, let μ_p be a Jensen measure for p on $A \cup B$. Next set

$$\lambda_0 = -\text{Im } z_1 + (\text{Im } z_1)^2 + (\text{Im } z_2)^2 - \frac{1}{2}(\text{Re } z_1)^2 - \frac{1}{2}(\text{Re } z_2)^2,$$

λ_0 is a strictly plurisubharmonic function, $\lambda_0(0) = 0$, and $\lambda_0 < -\frac{r^2}{2}$ on B . Therefore at least for p close to 0, one must have $\mu_p(A) > 0$. Hence if $\lambda = -\infty$ on A , one has for $p \in U$ close

to 0, $\lambda(p) = -\infty$, and therefore $\lambda = -\infty$ on any connected set containing 0 on which it is defined.

If one wishes to avoid using Jensen measures, one can argue very simply as follows: Fix $p \in U$ such that $\lambda_0(p) > -\frac{r^2}{4}$, if as above $\lambda = -\infty$ on A , then for $\epsilon > 0$ small enough $\lambda_0 + \epsilon\lambda$ clearly violates the maximum principle that in our case is $\text{Sup}_U = \text{Sup}_{A \cup B}$, unless $\lambda(p) = -\infty$.

The application to almost complex structures in real dimension 4 is immediate. Sharper statements are immediate too, and smoothness requirements are easy to lower (however not to $\mathcal{C}^1$, that would be interesting).

Proposition 4. *Let (M, J) be a connected almost complex manifold of class $\mathcal{C}^{3,\alpha}$ ($0 < \alpha < 1$), of real dimension 4. Let $p_0 \in M$. Let N be a germ of $\mathcal{C}^{3,\alpha}$ 2 dimensional surface, $p_0 \in N$. Assume that $J(p_0)TN_{p_0} \neq TN_{p_0}$, where TN_{p_0} is the tangent space to N at T_0 . Let λ be a J -plurisubharmonic function defined on M . If $\lambda = -\infty$ on N , then $\lambda \equiv -\infty$.*

Sketch of proof: It is enough to prove that $\lambda = -\infty$ on some non empty open set. We can then work in a neighborhood of p_0 . We can assume that $M \subset \mathbf{C}^2$, $p_0 = 0$, $J(0) = J_{st}$ (the standard complex structure on $\mathbf{C}^2$) and, after change of variables, that N coincides with $\mathbf{R}^2$. This change of variables may drop the regularity of J to $\mathcal{C}^{2,\alpha}$. Furthermore using dilations, as usual, one can assume that J is arbitrarily close in $\mathcal{C}^2$ norm to the standard complex structure. In that case, the function λ_0 considered above is J -plurisubharmonic, due to the characterization of plurisubharmonicity in terms of a Levi form (see e.g. [I-R] Corollary 1.1). Also, with U as above, C keeps its strong pseudoconcavity property. So the reasoning used in the complex case applies immediately if one proves that no non constant J -plurisubharmonic function on a neighborhood of U can attain its maximum on C . This follows from the fact that if $q \in C$ there is a J -holomorphic disc $u : \mathbf{D} \rightarrow U$, such that $u(0) = q$ and $u(\zeta) \in U$, if $\zeta \neq 0$ (see [I-R] Proposition 5.1).

3) Real dimension 2n. Of course non J -holomorphic surfaces can be J -pluripolar. Indeed $\mathbf{R}^2 \times \{0\}$ is pluripolar in $\mathbf{C}^3$ since it is included in the $-\infty$ set of $\log |z_3|$. The result here is that generic n -dimensional manifolds N ($TN \cup J(TN) = TM$) are not J -pluripolar. We just indicate how to adapt the reasoning for $\mathbf{C}^2$ above to the $\mathbf{C}^n$ case. Define U by:

$$\text{Im } z_1 > 0, \quad \sum_{j=2}^n (\text{Im } z_j)^2 < (\text{Im } z_1)^4, \quad \sum_{j=1}^n |z_j|^2 < r^2.$$

If again C is the part of the boundary of U corresponding to $\sum_{j=2}^n (\text{Im } z_j)^2 = \text{Im } z_1$, then C is of course not strictly pseudo-concave but at every point its Levi form has at least one negative eigenvalue, and this is all what is needed.

Appendix. Three Lemmas in Measure Theory.

Recall that we say that a subset $E \subset \mathbf{C}$ (on which the variable will now be denoted by z) is *fat* if 0 is a point of Lebesgue density of E , i.e. $\frac{|E \cap \{|z| < r\}|}{\pi r^2} \rightarrow 1$ as $r \rightarrow 0$, (where, the Lebesgue measure of a set $F \subset \mathbf{C}$ is denoted by $|F|$).

Lemma A1. *Let ν be a finite positive measure on $\mathbf{C}$. Assume that $\nu(\{0\}) = 0$, then there is a fat set E such that $z(\frac{1}{|z|} * \nu)$ tends to 0 as $z \rightarrow 0$, $z \in E$.*

Lemma A2. *Let $\psi \in L^1$ (on $\mathbf{C}$). Then, there is a fat set E such that $z^2(\psi * \frac{1}{z^2})$ tends to 0 as $z \rightarrow 0$, $z \in E$.*

The convolution makes sense by the theory of singular integrals.

Lemma A3. *Let $\rho \in L^1$, and $\epsilon > 0$, then the inequality $\rho > \frac{\epsilon}{|z|^2 |\log |z||}$ cannot be satisfied everywhere on a set F that has positive density at 0, i.e. satisfying $|F \cap \{|z| < r\}| > \delta r^2$ for some fixed $\delta > 0$, for all $r > 0$ small enough.*

Remarks.

1) It is enough to prove Lemma A3 for $\epsilon = 1$, replacing ρ by $\frac{\rho}{\epsilon}$. Lemma A3 will be enough for us, but obviously much stronger statements can be given. However, there may not exist any fat set on which $(|z|^2 |\log |z|| \rho)$ is bounded. An example is the following one. Take a sequence of integers (n_j) such that $\sum_j \frac{1}{\sqrt{n_j}} < +\infty$. For $2^{-n_j-1} < |z| < 2^{-n_j}$, set $\rho(z) = \frac{1}{|z|^2 \sqrt{|\log |z||}}$. Elsewhere set $\rho = 0$.

2) Simple logic shows that for proving Lemma A1, it is enough to prove that the set $E_\epsilon = \{z, \frac{1}{|z|} * \nu \leq \frac{\epsilon}{|z|}\}$ is a fat set, and again it is enough to prove this for $\epsilon = 1$. Then we can define E as follows. For a sequence $r_j > 0$ decreasing fast enough to 0 (in particular we require $r_{j+1} \leq \frac{1}{2}r_j$), define E by setting $E \cap \{r_{j+1} < |z| \leq r_j\} = E_{\frac{1}{j}} \cap \{r_{j+1} < |z| \leq r_j\}$.

By the same kind of arguments, for establishing Lemma A2, it is enough to prove that the set $E_1 = \{z, |\psi * \frac{1}{z^2}| \leq \frac{1}{|z|^2}\}$ is a fat set.

Proof of Lemma A1. As said above, we only need to prove that there is a fat set E such that $|z(\frac{1}{|z|} * \nu)| \leq 1$ on E . By writing ν as the sum of a measure with support near 0 and a measure with no mass near zero, the second one playing no role, we can assume $\|\nu\| \leq 1$. One has

$$\frac{1}{|z|} * \nu(z) = \int \int_{|z-t| \geq 2|z|} \frac{1}{|z-t|} d\nu(t) + \int_{|z-t| < 2|z|} \frac{1}{|z-t|} d\nu(t).$$

Trivially $\int \int_{|z-t| \geq 2|z|} \frac{1}{|z-t|} d\nu(t) \leq \frac{1}{2|z|}$. Fix $\epsilon > 0$, we need to prove that for $R > 0$ small enough, the set of z satisfying $|z| \leq R$ and where $\int_{|z-t| < 2|z|} \frac{1}{|z-t|} d\nu(t) \geq \frac{1}{2|z|}$ has measure at most ϵR^2 . For $R > 0$, let χ_R be the characteristic function of the set $|z| \leq R$, in $\mathbf{C}$. If $|z| \leq R$,

$$\int \int_{|z-t| < 2|z|} \frac{1}{|z-t|} d\nu(t) \leq A(z) = \left(\frac{1}{|\cdot|} \chi_{2R}(\cdot)\right) * (\chi_{3R} \nu)(z).$$

The standard estimates for L^1 norms of convolutions gives us

$$\|A\|_{L^1} \leq \left\| \frac{1}{|\cdot|} \chi_{2R} \right\|_{L^1} \|\chi_{3R}\nu\| = 4\pi R \|\chi_{3R}\nu\| \leq \frac{\epsilon}{2} R ,$$

if R is small enough (since ν has no point mass at 0). Then, obviously on the set $|z| \leq R$, $|A(z)| \geq \frac{1}{2|z|}$ can occur only on a set of measure at most ϵR^2 .

Proof of Lemma A2. Set $M = \{z, |\psi * \frac{1}{z^2}| > \frac{1}{|z|^2}\}$. It is enough to prove that, for any fixed $\delta > 0$, for $k \in \mathbf{N}$ large enough $|M \cap \{2^{-k-1} < |z| < 2^{-k}\}| \leq \delta 2^{-2k}$. Similarly to above we can write ψ as the sum of a function supported near 0, of small L^1 norm and a function vanishing near 0 and that plays no role. So we will be able to assume that the L^1 norm of ψ is as small as we will need. By the weak L^1 estimates in the Calderon Zygmund theory, we have for some universal constant C , for all $t > 0$:

$$|\{z \in \mathbf{C}, |\psi * \frac{1}{z^2}| > t\}| \leq C \|\psi\|_{L^1} \frac{1}{t}.$$

For $2^{-k-1} \leq |z| \leq 2^{-k}$, $|\psi * \frac{1}{z^2}| > \frac{1}{|z|^2}$ implies $|\psi * \frac{1}{z^2}| > 2^{2k}$, so the set of z satisfying $2^{-k-1} \leq |z| \leq 2^{-k}$ and $|\psi * \frac{1}{z^2}| > \frac{1}{|z|^2}$ has measure at most $\leq C \|\psi\|_{L^1} 2^{-2k}$. So it is enough to have $\|\psi\|_{L^1} \leq \frac{1}{C} \delta$, which we can assume, as said above.

Proof of Lemma A3.

Assume that $\rho > \frac{1}{|z|^2 |\log |z||}$ on a set F satisfying $|F \cap \{|z| < r\}| > \delta r^2$, for r small enough. Set $\eta = \sqrt{\frac{\delta}{2\pi}}$ (< 1). Then

$$|F \cap \{\eta r < |z| < r\}| > \frac{\delta}{2} r^2 .$$

Taking $r = \eta^k$ (k large), we get

$$\int_{F \cap \{\eta^{k+1} r < |z| < \eta^k\}} \rho > \frac{\delta}{2 |\log \eta|} \frac{1}{k} ,$$

clearly contradicting integrability of ρ , by summation over k .

REFERENCES.

- [D-S] K. Diederich, A. Sukhov. Plurisubharmonic functions and almost complex Stein structures. Mich. Math. J. (to appear)

- [E] F. Elkhadhra. J-pluripolar subsets and currents on almost complex manifolds. To appear in Math. Zeit.
- [Ga] T. Gamelin. *Uniform algebras*. Prentice Hall 1969.
- [G-S] H. Gaussier, A. Sukhov. Estimates of the Kobayashi metric on almost complex manifolds. Bull. Soc. Math. France 133 (2005) no 2, 259-273.
- [I-R] S. Ivashkovich, J-P. Rosay. Schwarz-type lemmas for solutions of $\bar{\partial}$ -inequalities and complete hyperbolicity of almost complex manifolds. Ann. Inst. Fourier 54 (2004), 2387-2435.
- [McD-S1] D. McDuff - D. Salamon. *J-holomorphic curves and quantum cohomology*. Univ. Lect. Series AMS, 6 (1994).
- [McD-S2] D. McDuff - D. Salamon. *J-holomorphic curves and symplectic topology*. AMS Colloquium Pub. 52 (2004).
- [N-W] A. Nijenhuis, W. Woolf. Some integration problems in almost complex and complex manifolds. Ann. of Math. **77** (1963), 424-489.
- [Pa] N. Pali. Fonctions plurisousharmoniques et courants positifs de type (1,1) sur une variété presque complexe. Manuscripta Math. 118 (2005) no 3, 311-337.
- [Ro1] J-P. Rosay. J-Holomorphic submanifolds are pluripolar. Math. Z. 253 (2006), 659-665.
- [Ro2] J-P. Rosay. Notes on the Diederich-Sukhov-Tumanov normalization for almost complex structures. Collect. Math. To appear.
Preprint: www.mittag-leffler.se/preprints/0708s/info.php?id=08
- [Si] J.-C. Sikorav. Some properties of holomorphic curves in almost complex manifolds. In *Holomorphic Curves in Symplectic Geometry*, eds. M. Audin and J. Lafontaine, Birkhauser, 1994, 351-361.
- [St] E.L. Stout. *Polynomial Convexity*. Progress in Math. 261. Birkhäuser 2007.
- [S-T1] A. Sukhov, A. Tumanov. Filling hypersurfaces by discs in almost complex manifolds of dimension 2. Indiana Univ. Math. J. **57** (2008), 509-544.
- [S-T2] A. Sukhov, A. Tumanov. Filling real hypersurfaces by pseudoholomorphic discs. J. Geom. Anal **18** (2008), 632-649.

Jean-Pierre Rosay
 Department of Mathematics
 University of Wisconsin
 Madison WI 53706, U.S.A.
jrosay@math.wisc.edu